

## References

1. Kovalevskaya G, Birken S, Kakuma T, Ozaki N, Sauer M, Lindheim S, et al. Differential expression of human chorionic gonadotropin (hCG) glycosylation isoforms in failing and continuing pregnancies: preliminary characterization of the hyperglycosylated hCG epitope. *J Endocrinol* 2002;172:497–506.
2. Elliott MM, Kardana A, Lustbader JW, Cole LA. Carbohydrate and peptide structure of the  $\alpha$ - and  $\beta$ -subunits of human chorionic gonadotropin from normal and aberrant pregnancy and choriocarcinoma. *Endocrine* 1997;7:15–32.
3. Frendo JL, Vidaud M, Guibourdenche J, Luton D, Muller F, Bellet D, et al. Defect of villous cytotrophoblast differentiation in syncytiotrophoblast in Down's syndrome. *J Clin Endocrinol Metab* 2000;85:3700–7.
4. Massin N, Frendo JL, Guibourdenche J, Luton D, Giovangrandi Y, Muller F, et al. Defect of syncytiotrophoblast formation and human chorionic gonadotropin expression in Down's syndrome. *Placenta* 2001;22(Suppl A):S93–7.
5. Cole LA, Shahabi S, Oz UA, Bahado-Singh RO, Mahoney MJ. Hyperglycosylated human chorionic gonadotropin (invasive trophoblast antigen) immunoassay: a new basis for gestational Down syndrome screening. *Clin Chem* 1999;45:2109–19.
6. Kovalevskaya G, Genbacev O, Fisher S, Caceres E, O'Connor J. Trophoblast origin of hCG isoforms: cytotrophoblasts are the primary source of choriocarcinoma-like hCG. *Mol Cell Endocrinol* 2002;194:147–55.
7. Birken S, Krichevsky A, O'Connor J, Schlatterer J, Cole L, Kardana A, et al. Development and characterization of antibodies to a nicked and hyperglycosylated form of hCG from a choriocarcinoma patient. *Endocrine* 1999;10:137–44.
8. Kovalevskaya G, Birken S, Kakuma T, O'Connor JF. Early pregnancy human chorionic gonadotropin (hCG) isoforms measured by an immunometric assay for choriocarcinoma-like hCG. *J Endocrinol* 1999;161:99–106.
9. Cole LA, Omrani A, Cermik D, Bahado-Singh RO, Mahoney MJ. Hyperglycosylated hCG, a potential alternative to hCG in Down syndrome screening. *Prenat Diagn* 1998;18:926–33.
10. Cole LA, Shahabi S, Oz UA, Rinne KM, Omrani A, Bahado-Singh RO, et al. Urinary screening tests for fetal Down syndrome: II. Hyperglycosylated hCG. *Prenat Diagn* 1999;19:351–9.
11. Bahado-Singh R, Oz U, Shahabi S, Omrani A, Mahoney M, Cole L. Urine hyperglycosylated hCG plus ultrasound biometry for detection of Down syndrome in the second trimester in a high-risk population. *Obstet Gynecol* 2000;95:889–94.
12. Khanlian SA, Smith HO, Cole LA. Persistent low levels of hCG: a pre-malignant gestational trophoblastic disease. *Am J Obstet Gynecol*; in press.
13. Diamandis EP, Christopoulos TK. The biotin-(strept)avidin system: principles and applications in biotechnology [Review]. *Clin Chem* 1991;37:625–36.
14. Weeks I, Beheshti I, McCapra F, Campbell AK, Woodhead JS. Acridinium esters as high-specific-activity labels in immunoassay. *Clin Chem* 1983;29:1474–9.
15. O'Connor JF, Ellish N, Kakuma T, Schlatterer J, Kovalevskaya G. Differential urinary gonadotropin profiles in early pregnancy and early pregnancy loss. *Prenat Diagn* 1998;18:1232–40.
16. Butler SA, Khanlian SA, Cole LA. Detection of early pregnancy forms of human chorionic gonadotropin by home pregnancy test devices. *Clin Chem* 2001;47:2131–6.

**Evaluation and Validation of a Duck IgY Antibody-based Immunoassay for High-Sensitivity C-reactive Protein: Avian Antibody Application in Clinical Diagnostics**, Yun-Chien Tsen,<sup>1</sup> Guor-Yuan Kao,<sup>1</sup> Chung-Li Chang,<sup>1</sup> Fang-Yu Lai,<sup>1</sup> Chien-Hsun Huang,<sup>1</sup> San Ouyang,<sup>1</sup> Mei-Huei Yu,<sup>2</sup> Chi-Pin Wang,<sup>3</sup> and Y-Neng Chiou<sup>1\*</sup> (<sup>1</sup> Good Biotech Corp., No. 38 34th Road, Taichung Industrial Park, Taichung 407, Taiwan; <sup>2</sup> Kuang Tien General Hospital Laboratory, Taichung 433, Taiwan; <sup>3</sup> Chung Shan Medical University Hospital, Taichung 402, Taiwan; \* author for correspondence: fax 886-4-23590410, e-mail gbc@good-biotech.com)

High-sensitivity C-reactive protein (hs-CRP) is considered one of the most promising markers for cardiac risk assessment (1). A highly sensitive and interference-resistant CRP assay plays an important role in accurate CRP

measurements (2, 3); therefore, manufacturers of in vitro diagnostics have begun modifying their existing CRP assays to meet the requirements. Mammalian antibodies, e.g., rabbit, mouse, and goat IgGs, are the major ones used in clinical immunoassays, but they have some drawbacks. Within the past 10 years, it has been reported that rheumatoid factor and heterophilic antibodies present in samples have caused false-positive results in CRP measurements (4–6). In addition, monoclonal antibodies for use as diagnostic and therapeutic reagents can induce human anti-mouse antibodies, which may interfere with mammalian IgGs in immunoassays, especially in the sandwich-type methods (7).

To eliminate interference without increasing costs, avian antibodies are now considered potential candidates to improve current immunoassays. One such assay, the CRPex-HS<sup>®</sup> C-Reactive Protein LIT assay (Good Biotech Corp., Taiwan), based on latex microparticle-enhanced immunoturbidimetry, has been developed and is commercially available. The CRPex-HS CRP LIT assay is intended to measure serum hs-CRP concentrations with use of a duck antibody. The duck anti-CRP IgY ( $\Delta$ Fc) antibody is a good substitute for traditional mammalian antibodies. The antibody, purified from egg yolks of white Pekin ducks (*Anas platyrhynchos*), is coupled to polystyrene microparticles. When CRP in the serum sample encounters microparticles containing the duck anti-CRP IgY ( $\Delta$ Fc), it agglutinates and the turbidity is measured at 570 nm. The assay calibration range is 0–20.0 mg/L. Practical CRP measurements by the CRPex-HS CRP LIT assay are conducted on a common automated chemistry analyzer, the Hitachi 717. In this study, we evaluated and validated the performance characteristics of this new hs-CRP assay.

The specificity of the duck anti-CRP IgY ( $\Delta$ Fc) used in the hs-CRP assay was assessed by Western blot analysis. Both CRP in the patient serum and the partially purified CRP (Crystal Chem Inc.) were specifically recognized by the duck anti-CRP IgY ( $\Delta$ Fc).

To study linearity, we diluted a patient serum pool with a CRP concentration of 20.0 mg/L with CRP-free serum to the following final percentages of the patient pool: 100%, 75%, 50%, 25%, 12.5%, 6.25%, and 0.5%. The samples were analyzed in triplicate in a single analytical run. The linearity of the method was confirmed up to 20 mg/L (i.e., the highest calibrator concentration). Regression analysis of the data yielded an intercept of  $-0.12$  mg/L, a slope of 21.69, and an  $r^2$  of 1.0. One control (Cliniq Corporation) with a very high CRP concentration was diluted to test the possibility of underestimating the true CRP value because of a prozone effect. We found that a CRP concentration of up to 500 mg/L did not cause a false-negative result. Concentrations  $>500$  mg/L were not tested. The detection limit, 0.05 mg/L CRP, was the concentration corresponding to a signal 2 SD above the mean for a CRP-free serum.

Within-run imprecision (as CVs), determined by analyzing three serum samples (mean values, 0.16, 5.56, and 18.65 mg/L) in 20 replicates, were 10%, 0.73% and 0.98%, respectively. For day-to-day imprecision, samples were




Fig. 1. Comparison of CRP concentrations determined by the IgY-based method with other commercial assays.

(A–C), regression analysis; (D–F), Bland–Altman plots. ○, samples from control individuals; ×, samples from avian-allergic individuals. In panels A–C, the solid lines are the regression lines; in panels D–F, the solid lines are the means, and the dashed lines are  $\pm 2$  SD.

analyzed daily in triplicates, and the daily averages for a total of 14 days were calculated. CVs were 5.2%, 2.5%, and 3.9% at (mean) CRP concentrations of 0.37, 4.02, and 12.91 mg/L. The functional sensitivity, the lowest measurable CRP concentration with a CV  $<20\%$ , was 0.058 mg/L. Interference was defined as a difference of  $>5\%$  between the assay values of the original samples and those of samples to which various concentrations of interferent had been added. No interference occurred with up to 450

mg/L unconjugated bilirubin (cat. no. B-4126; Sigma). Hemoglobin (prepared from fresh hemolysate) up to 9400 mg/L and fat emulsion (Liposyn® II; Abbott Laboratories) up to 10 g/L also did not interfere with the CRP assay.

Because the source of duck IgY ( $\Delta$ Fc) is the duck egg and an allergic reaction to eggs is generally agreed to be one of the most common causes of food allergies in infants and young children, it is likely that IgE antibodies against duck antibodies will be induced in avian-allergic patients

**Table 1. Comparison of samples from the three CRP particle-enhanced immunoturbidimetric assays for control and avian-allergic individuals.**

$y$ -Axis method	$x$ -Axis method	Slope	$y$ -Intercept, mg/L	$R^2$	Samples	Fig. 1 panel
CRPex-HS	Wako CRP-UL	1.0875	0.0022	0.9981	Control	A
CRPex-HS	Wako CRP-UL	1.0083	0.0587	0.9935	Avian-allergic	A
CRPex-HS	K-ASSAY	0.9744	-0.0376	0.9982	Control	B
CRPex-HS	K-ASSAY	0.9678	0.0008	0.9957	Avian-allergic	B
K-ASSAY	WAKO CRP-UL	1.1154	0.0422	0.9986	Control	C
K-ASSAY	Wako CRP-UL	1.0398	0.0623	0.9940	Avian-allergic	C

during their lifetimes. One study regarding egg-allergic patients that demonstrated specific IgE binding against egg yolk-derived antiviral chicken immunoglobulins (8) prompted us to assess potential interference from the serum of avian-sensitized individuals in the duck antibody-based immunoassay. Two commercial CRP assays, the Wako CRP-UL CRP (Wako Pure Chemical Industries, Ltd.) and the K-ASSAY<sup>®</sup> CRP (2) (Kamiya Biomedical Company) assays, which use goat IgG and rabbit IgG, respectively, were used to assay the samples on a Hitachi 717 analyzer. The results were interpreted by least-squares regression analysis and Bland–Altman plots.

We obtained 76 serum samples from individuals shown to have IgE antibodies to egg yolks, egg whites, feathers, and other avian products or byproducts. As controls, the remaining serum samples from 86 individuals who had undergone a physical checkup were collected consecutively from local hospitals. The use of remaining sera for evaluation of *in vitro* diagnostic methods was in concordance with local regulations. To ensure the privacy of the human research subjects, neither the names nor the identification numbers linked to the samples were provided. Serum samples of allergic individuals were randomly interspersed between the control samples and were analyzed simultaneously with the Wako CRP-UL CRP, K-ASSAY CRP (2), and the CRPex-HS CRP LIT assays. All three CRP assays were used according to the manufacturers' instructions.

Comparison of the CRPex-HS vs Wako CRP-UL assays revealed a slight difference between the regression line for samples from control individuals and that for avian-allergic samples (Fig. 1A). However, comparison of the CRPex-HS and the K-ASSAY CRP (2) showed a less-pronounced difference between the regression lines (Fig. 1B). The noticeable difference might be attributable to the divergent distribution of CRP values in the two sample populations. The calculated regression equations are compared in Table 1.

From the Bland–Altman plots, we identified two samples for which the IgY ( $\Delta$ Fc)-based method seemed to produce incorrect CRP results (Fig. 1, D and E, arrows), but we were unable to identify the source of the problem. According to the proposed algorithm for cardiovascular risk assessment using CRP (9), risk stratification of patients, regardless of the assay used, was not mismatched by more than one adjacent risk group. Generally, the IgY ( $\Delta$ Fc)-based method correlated well with other mamma-

lian IgG-based assays for CRP quantification of the general population, including avian-allergic individuals.

Duck IgY ( $\Delta$ Fc), compared with chicken IgY, lacks two C-terminal domains (C $\nu$ 3, C $\nu$ 4) of the heavy chains and is structurally equivalent to the F(ab')<sub>2</sub> fragment (10). Ducks, as with chickens, pass immunity to their offspring by transferring antibodies to eggs; we therefore isolated the specific duck IgY ( $\Delta$ Fc) from the collected yolks. Avian antibody technology is a humane and cost-effective method of antibody production. In addition, avian IgY has been shown to have poor cross-reactivity to mammalian IgG, Fc receptor, complement components, and rheumatoid factor because of structural differences with mammalian IgG antibody, thus eliminating false-positive results and human anti-mouse antibody interference (11–14). One study showed that the largely phylogenetic difference between avian host animals and the antigens of mammalian sources increases the immune response; thus, avian IgY recognized more epitopes than the corresponding mammalian IgG and was relatively more reactive to the highly conserved proteins (15). It has been proposed that more widespread use of avian antibodies be promoted (16).

In the past, difficulties in separating antibodies from macromolecules, lipids, and many proteins in duck egg yolks hampered the popularization of duck antibodies for clinical and research purposes. Our study demonstrates that the duck antibody-based CRP assay is comparable to mammalian antibody-based assays. Duck IgY ( $\Delta$ Fc) antibody can be regarded not only as a multipurpose tool in laboratory research but also an alternative candidate for diagnostic applications.

## References

- Rifai N, Ridker PM. High-sensitivity C-reactive protein: a novel and promising marker of coronary heart disease [Review]. *Clin Chem* 2001;47:403–11.
- Roberts WL, Moulton L, Law TC, Farrow G, Cooper-Anderson M, Savory J, et al. Evaluation of nine automated high-sensitivity C-reactive protein methods: implications for clinical and epidemiological applications. Part 2. *Clin Chem* 2001;47:418–25.
- Ohtake T, Kano S, Watanabe K. Interference in turbidimetric immunoassay for serum C-reactive protein due to serum protein abnormalities an immune complex and rheumatoid factor. *Rinsho Byori* 2000;48:752–9.
- Muller W, Mierau R, Wohltmann D. Interference of IgM rheumatoid factor with nephelometric C-reactive protein determinations. *J Immunol Methods* 1985;80:77–90.
- Tsuda H, Takata T, Nagafuchi S, Etoh F, Kurihara M, Okochi K, et al. Falsely increased nephelometric results caused by a monoclonal rheumatoid factor. *Clin Chem* 1990;36:1263–4.
- Benoist JF, Orbach D, Biou D. False increase in C-reactive protein attribut-

- able to heterophilic antibodies in two renal transplant patients treated with rabbit antilymphocyte globulin. *Clin Chem* 1998;44:1980–5.
7. Klee GG. Human anti-mouse antibodies. *Arch Pathol Lab Med* 2000;124:921–3.
  8. Bernhisel-Broadbent J, Yolken RH, Sampson HA. Allergenicity of orally administered immunoglobulin preparations in food-allergic children. *Pediatrics* 1991;87:208–14.
  9. Rifai N, Ridker PM. Proposed cardiovascular risk assessment algorithm using high-sensitivity C-reactive protein and lipid screening. *Clin Chem* 2001;47:28–30.
  10. Warr GW, Magor KE, Higgins DA. IgY: clues to the origins of modern antibodies [Review]. *Immunol Today* 1995;16:392–8.
  11. Lindahl TL, Festin R, Larsson A. Studies of fibrinogen binding to platelets by flow cytometry: an improved method for studies of platelet activation. *Thromb Haemost* 1992;68:221–5.
  12. Larsson A, Wejaker PE, Forsberg PO, Lindahl T. Chicken antibodies: a tool to avoid interference by complement activation in ELISA. *J Immunol Methods* 1992;156:79–83.
  13. Larsson A, Karlsson-Parra A, Sjöquist J. Use of chicken antibodies in enzyme immunoassays to avoid interference by rheumatoid factors. *Clin Chem* 1991;37:411–4.
  14. Larsson A, Mellstedt H. Chicken antibodies: a tool to avoid interference by human anti-mouse antibodies in ELISA after in vivo treatment with murine monoclonal antibodies. *Hybridoma* 1992;11:33–9.
  15. Gassmann M, Thommes P, Weiser T, Hubscher U. Efficient production of chicken egg yolk antibodies against a conserved mammalian protein. *FASEB J* 1990;4:2528–32.
  16. Schade R, Staak C, Hendriksen C, Erhard M, Hugl H, Koch G, et al. The production of avian (egg yolk) antibodies: IgY. The report and recommendations of ECVAM Workshop 21. *Altern Lab Anim* 1996;24:925–34.

**Influence of Sampling-Time Error on Cyclosporine Measurements Nominally at 2 Hours after Administration**, Frank Saint-Marcoux,<sup>1</sup> Annick Rousseau,<sup>1,2</sup> Yann Le Meur,<sup>3</sup> Marc Estenne,<sup>4</sup> Christiane Knoop,<sup>4</sup> Jean Debord,<sup>1</sup> and Pierre Marquet<sup>1\*</sup> (Departments of <sup>1</sup>Pharmacology and Toxicology and <sup>3</sup>Nephrology, University Hospital, 87042 Limoges cedex, France; <sup>2</sup>Laboratory of Biophysics, Faculty of Pharmacy, 87025 Limoges cedex, France; <sup>4</sup>Departments of Chest Medicine and Clinical Chemistry, Erasme University Hospital, 1070 Brussels, Belgium; \* address correspondence to this author at: Service de Pharmacologie et Toxicologie, CHU Dupuytren, 87042 Limoges cedex, France; fax 33-555-05-61-62, e-mail marquet@unilim.fr)

Cyclosporine (CsA) blood concentrations measured 2 h after Neoral<sup>®</sup> administration ( $c_2$ ) are a sensitive predictor of clinical outcome in organ transplantation, as suggested by a recent prospective clinical trial in liver transplant patients (1).  $c_2$  is now recommended as the target exposure index for the therapeutic drug monitoring (TDM) of CsA (2–8). The aim of this study was to investigate, for different types of grafts, the concentration–time relationships around  $c_2$  to evaluate the concentration error as a function of the sampling-time error and to identify the sampling-time range compatible with acceptable performance of this  $c_2$  TDM strategy.

Data obtained from three different clinical trials were studied retrospectively. Each patient gave written informed consent, and each trial was approved by a local ethics committee (9–11). The three populations were as follows:

- Twenty stable adult renal transplant patients (grafted for more than 3 months and with stable renal function), from whom samples were collected immediately before a morning dose ( $t_0$ ) and at 0.33, 0.66, 1, 1.5, 2, 3, 4, 6, and 9 h post dose. These 20 full profiles were used for this study.
- Sixteen heart transplant patients for whom three full pharmacokinetic profiles (12 samples) were obtained, at 1 week ( $W_1$ ), 12 weeks ( $M_3$ ), and 52 weeks ( $Y_1$ ) after transplantation. Samples were collected immediately before and at 0.33, 0.66, 1, 1.5, 2, 2.5, 3, 4, 6, 9, and 12 h after a morning dose. Sixteen profiles were available at  $W_1$ , and 29 were available for  $M_3$  and  $Y_1$  combined. We previously found no statistical difference between these two time periods with respect to pharmacokinetic parameters, whereas there were significant differences between  $W_1$  and either  $M_3$  or  $Y_1$ . For this reason the last two periods were pooled (11).
- Twenty stable adult lung transplant patients with ( $n = 10$ ) and without ( $n = 10$ ) cystic fibrosis. Three full pharmacokinetic profiles were collected within 1 week for each patient at 0, 0.5, 1, 1.5, 2, 2.5, 3, 4, 5, 6, 8, and 12 h after a morning dose. A total of 29 profiles were obtained from cystic fibrosis transplant patients, and 30 from non-cystic fibrosis patients.

All the patients from these three clinical trials were dosed twice daily with microemulsified CsA (Neoral). The data are summarized in Table 1.

For both kidney and lung transplant patients, CsA whole-blood concentrations were measured using an enzyme-multiplied immunoassay technique (Emit; Dade-Behring Diagnostics). This method has an upper limit of quantification of 500  $\mu\text{g/L}$ . In the clinical trial conducted with cardiac transplant recipients, whole-blood CsA was measured with a fluorescence polarization immunoassay (FPIA; Abbott TDx). This method has an upper limit of quantification of 1500  $\mu\text{g/L}$ . For both assays, samples with CsA concentrations greater than the upper assay range were diluted 1:4 with human blank whole blood (100  $\mu\text{L}$  of sample + 300  $\mu\text{L}$  of blank whole blood) and then reanalyzed.

Individual pharmacokinetic profiles were fitted using

**Table 1. Mean CsA exposure indices in the populations studied.**

Type of graft	No. of full profiles	Mean (SD) measured $c_0$ , $\mu\text{g/L}$	Mean (SD) measured $c_2$ , $\mu\text{g/L}$
Kidney			
Stable (>3 months)	20	134.6 (41.3)	775.6 (188)
Lung			
Cystic fibrosis	29	267.9 (91.1)	1453 (667)
Non-cystic fibrosis	30	318.5 (97.2)	1472 (429)
Heart			
De novo (week 1)	16	329.3 (119)	1010 (311)
Stable (month 3 and year 1)	29	287 (99)	1242 (318)